

## Amplite™ Colorimetric Aldehyde Quantitation Kit

Ordering Information	Storage Conditions	Instrument Platform
Product Number: 10051 (200 assays)	Keep at -20 °C Avoid moisture and light	Absorbance microplate readers

### Introduction

Very reactive aldehydes, namely 4-hydroxyalkenals, were first shown to be formed in autoxidizing chemical systems. It was subsequently shown that 4-hydroxyalkenals, particularly 4-hydroxynonenal, were formed in substantial amounts under biological conditions, i.e. during the peroxidation of lipids of liver microsomes incubated in the NADPH-Fe system. Many other aldehydes were also identified in peroxidizing liver microsomes or hepatocytes, e.g., alkanals, alk-2-enals, and 4-hydroxyalkenals.

Most of the existing aldehyde test methods are based on separations either by the tedious and expensive HPLC-MS or GC-MS. Our Amplite™ Colorimetric Aldehyde Quantitation kit uses a proprietary dye that generates a chromogenic product upon reacting with an aldehyde. The kit provides a sensitive, one-step colorimetric method to detect as little as 1 nanomole of aldehyde in a 100 µL assay volume (10 µM). The assay can be performed in a convenient 96-well or 384-well microtiter-plate format and readily adapted to automation without a separation step. Its signal can be easily read with an absorbance microplate reader at 405 or 550 nm. This kit has been used for monitoring activities of oxidases that convert an amino group to an aldehyde group.

### Kit Key Features

<b>Broad Application:</b>	Can be used for quantifying aldehydes in a variety of applications such as carbohydrate, lipid chemistry, as well as enzyme reactions.
<b>Sensitive:</b>	Detect as low as 1 nanomole of aldehyde.
<b>Continuous:</b>	Easily adapted to automation without a separation step.
<b>Convenient:</b>	Formulated to have minimal hands-on time. No wash is required.
<b>Non-Radioactive:</b>	No special requirements for waste treatment.

### Kit Components

Components	Amount
Component A: AldeView™ Yellow	2 bottles
Component B: Assay Solution	1 bottle (10 mL)
Component C: Aldehyde Standard	1 vial
Component D: Dilution Buffer	1 bottle (20 mL)

### Assay Protocol for One 96-Well Plate

#### Brief Summary

**Prepare enzyme reaction (50 µL) → Add 2X AldeView™ Yellow reaction mixture (50 µL) → Incubate at room temperature for 30 to 60 minutes → Monitor absorbance increase at 405 or 550 nm**

*Note: Thaw all the kit components to room temperature before starting the experiment.*

#### 1. Prepare 2X AldeView™ Yellow reaction mixture:

Add 5 mL of Assay Solution (Component B) into the bottle of AldeView™ Yellow (Component A), and mix well.

*Note 1: 5 mL of the 2X AldeView™ Yellow reaction mixture is enough for 1 plate. The reaction mixture is not stable. Use within 2 hours.*


*Note 2: Assay solution (Component B) is potentially hazardous. Wear gloves when handling it.*



## Data Analysis

The absorbance in blank wells (with 0  $\mu\text{M}$  aldehyde standard and 2X AldeView™ Yellow reaction mixture only) is used as a control, and is subtracted from the values for those wells with the aldehyde reactions. An aldehyde standard curve is shown in Figure 1.

*Note: The absorbance background increases with time, thus it is important to subtract the absorbance intensity value of the blank wells for each data point.*



**Figure 1.** Aldehyde dose response was measured in a white/clear bottom 96-well plate with Amplitude™ Colorimetric Aldehyde Quantitation Assay Kit using a Spectrum Max microplate reader (Molecular Devices). As low as 10  $\mu\text{M}$  (1 nanomol/well) of aldehyde can be detected with 30 minutes incubation (n=3).

## References

1. Trevor M. Kitson. (1985) High concentrations of aldehydes slow the reaction of cytoplasmic aldehyde dehydrogenase with thiol-group modifiers Biochem. J. 228, 765.
2. Crabb DW, Matsumoto M, Chang D, You M (2004). Overview of the role of alcohol dehydrogenase and aldehyde dehydrogenase and their variants in the genesis of alcohol-related pathology. The Proceedings of the Nutrition Society 63 (1): 49.
3. Steinmetz CG, Xie P, Weiner H, Hurley TD (1997). Structure of mitochondrial aldehyde dehydrogenase: the genetic component of ethanol aversion. Structure 5 (5): 701.
4. O'Donnell JM, Kudej RK, LaNoue KF, Vatner SF, Lewandowski ED. (2004) Limited transfer of cytosolic NADH into mitochondria at high cardiac workload. Am J Physiol Heart Circ Physiol, 286, H2237.
5. Zurek G, and karst U (2000). 2,4-Dinitro-3,5,6-trideuterophenylhydrazones for the quantitation of aldehydes and ketones in air samples by liquid chromatography-mass spectrometry. J of chromatography A, 869, 251.
6. Ou Z, Ogamo A, Guo L, Konda Y, Harigaya Y, and Nakagawa Y. (1995). Identification and quantitation of choline glycerophospholipids that contain aldehyde residues by fluometric high-performance liquid chromatography. Analytical biochemistry 227, 289.

**Warning: This kit is only sold to end users. Neither resale nor transfer to a third party is allowed without written permission from AAT Bioquest. Chemical analysis of kit components is strictly prohibited. Please call us at 408-733-1055 or e-mail us at [info@aatbio.com](mailto:info@aatbio.com) if you have any questions.**